

Facts on Colorectal Cancer



Get tested for Colorectal Cancer

You have probably heard that early detection of cancer saves lives. This is particularly true for cancer of the colon and rectum, or colorectal cancer. Colorectal cancer is the 3rd most common cause of cancer worldwide and the 2nd leading cause of cancer deaths in the U.S.

The large intestine or colon is the lowest portion of the digestive system and measures about 5 feet long. The rectum is the last 6 inches of the colon. Colorectal cancer usually develops gradually over several years. It often begins as a benign (non-cancerous) polyp, a small growth of tissue in the lining or wall of the colon or rectum. Over time, some of these polyps can grow and become malignant (cancerous). As they grow, polyps can bleed or block the intestine. By performing early screening and removing all small visible benign polyps, colorectal cancer can be prevented.

Who gets colorectal cancer?

Persons with the following risk factors are more likely to develop colorectal cancer:

- Age
Over 90% of colorectal cancer occurs in persons older than 50.
- Family or personal history of colorectal cancer, polyps, or chronic inflammatory bowel disease
Note: Only 25% of persons diagnosed to have colorectal cancer have a positive family history of the disease. 75% of colorectal cancer occurs in persons without a family history.

- Physical inactivity
- Overweight or obesity
- Cigarette smoking
- Diet high in fat and meat
- Diet low in fiber, fruits, and vegetables
- High intake of alcohol

Warning signs of colorectal cancer

As with most cancers, early colorectal cancer frequently has no symptoms. As the cancer becomes larger, signs and symptoms may include:

- A change in bowel habits (diarrhea or constipation)
- A change in the shape of the stools (becomes narrower)
- Feeling that bowel does not empty completely
- Abdominal discomfort (bloating, cramping, pain)
- Blood in the stool (either bright red or dark in color)
- Bleeding from the rectum
- Chronic fatigue and weakness
- Unexplained weight loss

Colorectal screening tests and guidelines

Beginning at age 50, the American Cancer Society recommends that both men and women at average risk for developing colorectal cancer follow one of the screening tests listed below. Persons at higher risk because of family history or medical conditions should begin screening earlier and more often as recommended by their physician. Most screening tests, with the exception of fecal occult blood test (FOBT), require that laxatives be taken and enemas used before the exam to clean out the colon for better viewing.

1. Fecal occult blood test (FOBT) every year

This test can detect very small quantities of blood or hidden blood in the stool. Stool samples are collected and thinly spread onto special cards which are then returned to a medical lab for testing. A special diet may be required a few days before the test.

2. Flexible sigmoidoscopy every 5 years

The sigmoid colon (shaped like an S) is the end section of the colon, which leads to the rectum. A thin, hollow, lighted tube called a sigmoidoscope is inserted through the rectum and into the colon, allowing the doctor to see about one-third of the colon.

3. Colonoscopy every 10 years

An instrument, similar to a sigmoidoscope but longer, allows the doctor to see the entire colon and also remove polyps at the same time. Colonoscopy is the most sensitive method for detection of cancerous polyps.

4. Double-contrast barium enema every 5 years

A chalky substance called barium sulfate is used to fill the colon so that X-rays can be taken to detect polyps or abnormalities.

5. CT colonography (virtual colonoscopy) every 5 years

CT colonography uses low dose radiation CT scanning to obtain an inside view of the colon. CT scanning combines special x-ray equipment with computer software to create multiple images which can show polyps or abnormalities. Whether CT colonography can lower the number of deaths from colorectal cancer is unknown at this time.

Note: Any positive or abnormal test results should be followed by a colonoscopy for a more complete diagnosis.

Treatment for colorectal cancer

For cancer that is in the early stage, surgery is the most common form of treatment and often results in a very high cure rate.

For cancer that has spread to the nearby lymph nodes or other organs, chemotherapy (the use of drugs to kill cancer cells) or chemotherapy with radiation may be recommended before or after surgery.

Prevention of colorectal cancer

- Get screened for colorectal cancer
- Eat a plant-based diet with plenty of whole grains, fruits, and vegetables
- Reduce intake of high fat foods especially from animal sources
- Be physically active
- Achieve and maintain a healthy weight
- Limit alcohol consumption
- Stop smoking

Studies show that taking aspirin and postmenopausal hormones decrease the risk of colorectal cancer.



華人社區健康資源中心
Chinese Community Health Resource Center
www.cchrhealth.org

© 2006 CCHRC. Revised 2013.

Printing made possible through funding from National Cancer Institute for Asian American Network for Cancer Awareness, Research and Training - Community Network Program (AANCART-CNP) grant number U01 CA 114640